

REHAB QUIZ

- 1. Drug used in facet intra-articular facet joint injection is**
 - A) Bupivacaine 0.25%
 - B) Bupivacaine 0.5%
 - C) Bupivacaine 0.75%
 - D) Bupivacaine 1.25%

- 2. Percutaneous discectomy is indicated in**
 - A) PIVD extrusion
 - B) PIVD protrusion
 - C) PIVD sequestration
 - D) None

- 3. All diagnostic selective nerve root block is helpful in referred pain to chest except**
 - A) C6
 - B) C7
 - C) C8
 - D) T1

- 4. Intradiscal injection is indicated in coccydynia with**
 - A) Most painful area over tip
 - B) Spicule
 - C) Absence of radiologic lesion in dynamic film
 - D) Dynamic film confirmation of luxation or hypermobility.

- 5. All are contra-indications of intra-articular hyaluronic acid in OA knee except**
 - A) Ligamentous laxity
 - B) DM
 - C) Bleeding diatheses
 - D) Allergy (avion)

- 6. Which interventions are indicated if back and thigh pain more then leg pain in case of degenerative spondylolsthesis of L4/L5**
 - A) Transforaminal ESI
 - B) Caudal epidural
 - C) L4-5 facet joint block
 - D) All

7. Ozone therapy are indicated in all except

- A) Knee OA
- B) AVN of femoral head
- C) Multilevel PIVD
- D) Shoulder pathology

8. Most curative mode of intervention in facet joint arthropathy

- A) Laser
- B) IDET technique
- C) Radiofrequency ablation
- D) Vertebroplasty

9. Botulinium toxin A acts by

- A) Preventing release of Ach
- B) Preventing uptake of Ach
- C) Preventing release of Adrenaline
- D) Preventing uptake of Adrenaline

10. In a spastic paraplegic patient perineal hygiene is better maintained by phenol block of

- A) Femoral nerve
- B) Sciatic nerve
- C) Obturator nerve
- D) Post-tibial nerve.

**Please send the answer to the editor with your name, address of communication.
The right responder will be acknowledged in next issue of journal.**