## **PG Forum**

## **REHAB QUIZ**

- 1. During which phase of throwing is the elbow joint placed under the most valgus stress?
  - a) Follow through
  - b) Wind up
  - c) Early cocking
  - d) Late cocking
- 2. Which of the following does the nerve conduction component of the neurodiagnostic study fail to assess or give information about?
  - a) Autonomic nerve
  - b) Integrity of myelin
  - c) Sensory nerve
  - d) Speed of transmission
- 3. The X axis of oscilloscope (screen) represents.
  - a) Time in microseconds
  - b) Time in milliseconds
  - c) Distance in centimetres
  - d) Distance in millimetres
- 4. As opposed to acquired neuropathies, congenital neuropathies usually.
  - a) Have proximal more than distal slowing
  - b) Have uniform slowing throughout the nerve
  - c) Have segmental slowing throughout the nerve
  - d Are distal
- 5. Monopolar needles generally have higher amplitude motor that action potential (MUAPS) than concentric needles because.
  - a) Monopolar needles are longer than concentric needles
  - b) The needles samples from 360 degree rather than 180 degree
  - c) The tip of a concentric needle is smaller
  - d) Concentric needles have the ground electrode as part of the needle

- 6. All of the following are true about piriformis syndrome except
  - a) The sciatic nerve may be involved because in some individual the nerve releases through piriformis muscle
  - b) Pain may be in lat buttock, post hip to thigh, sciatica like pain
  - c) Pain in flexion, abduction, external rotation
  - d) Pain may be exacerbated by walking upstairs or prolong sitting

## 7. What is terrible triad?

- a) Medial/lateral meniscal injury with anterior cruciate ligament(ACL) tear
- ACL, medial collateral ligament(MCL),and medial meniscus injury
- c) ACL, post cruciate ligament(PCL),and lat meniscal injury
- d) ACL, MCL, and PCL
- 8. Mallet finger is.
  - a) A rupture of terminal extensor tendon of distal phalanx
  - b) Identified by loss of active extension of proximal interphalangeal joint(PIP) of finger
  - Caused by forced extension of distal phalangeal joint
  - d) Occurs more commonly in ice hockey than in baseball or basketball
- 9. Little league elbow.
  - a) Involves the lateral elbow region
  - b) Is an acute dislocation of elbow
  - c) Occurs most commonly b/w ages 3-15
  - d) Occurs in athletes complaining of medial elbow pain
- 10. An injury involving the centre of optic chiasm would result in.
  - a) Homonymus hemianopia
  - b) Bitemporal hemianopia
  - c) Cortical blindness
  - d) Monoocular blindness

## ANSWERS